FLORIDA CAC, CAP-TRAINING & SUPERVISION PROGRAM

(ADDICTION COUNSELOR EXAM PREP) COURSEBOOK

CICE RIVERA, MS, PHD, CAP

TABLE OF CONTENTS

Clinical Evaluation ……..……..…….05
Clinical Evaluation Continued ……...…….55
Counseling Fundamentals. ……..……89
Counseling & Treatment Planning ……..……122
Case Management and Referral ……..……154
Ethical Professional, Trauma and Intervention ……..…….……209
Ethics Code of Conduct, Policies, Crisis……..…….242
Co - Occurring Disorder, Treatment Knowledge ……..…….…….271
Practicum and Supervised Experience I ……..…… 330
Practicum and Supervised Experience II ……..……361
Client Session, Portfolio, Mentorship ……..390
Client Integration, Case, Certification…….……..……418
Capstone, Practice, Career…….……..…….…….. 447

INTRODUCTION TO ADDICTION AND COUNSELING FOUNDATIONS-DAY 1

Addiction is a chronic brain disorder characterized by compulsive substance use despite harmful consequences. It can involve substances such as alcohol, opioids, stimulants, cannabis, and prescription medications, as well as behavioral addictions like gambling or excessive internet use. Addiction affects the brain's reward system, influencing decision-making, impulse control, and the ability to experience pleasure naturally. Individuals may experience strong cravings, withdrawal symptoms, or repeated relapse, which makes understanding the underlying causes essential for effective intervention.

Recognizing addiction requires awareness of both behavioral and physical signs. Common indicators include neglect of responsibilities, social withdrawal, changes in mood, secrecy, and continued substance use despite negative consequences. Contributing factors can include genetic predisposition, family history, environmental stressors, trauma, and mental health conditions. Counselors must also consider age, developmental stage, and life circumstances when assessing clients to provide a comprehensive understanding of their substance use behaviors.

Human development significantly influences substance use patterns and treatment needs. Adolescents, whose brains are still developing, are highly susceptible to peer pressure and risk-taking behaviors. Adults may struggle with addiction in response to stressors like work demands, relationships, or mental health challenges. Older adults may face unique challenges, including health complications, social isolation, or polypharmacy. By understanding developmental stages, counselors can tailor interventions that are appropriate for the client's age, cognitive abilities, and life experiences, enhancing the likelihood of positive outcomes.

As a Certified Addiction Counselor and or Professional (CAC), (CAP), the counselor's responsibilities include conducting assessments, developing individualized treatment plans, implementing counseling interventions, and monitoring client progress. Maintaining ethical standards is critical, including confidentiality, accurate

documentation, and adherence to mandatory reporting laws. Professional boundaries and avoidance of dual relationships ensure that the counseling relationship remains therapeutic and effective. Awareness of legal and ethical obligations, combined with knowledge of addiction and human development, allows counselors to provide safe, effective, and client-centered care.

CLINICAL EVALUATION AND FOUNDATIONS

Clinical evaluation is the cornerstone of effective substance use counseling. It involves systematically gathering information about a client's physical, psychological, emotional, and social functioning to develop an accurate understanding of their needs. For both CACs and CAPs, clinical evaluation begins with establishing rapport and engaging clients in a collaborative process. Understanding the client's history, presenting concerns, and motivation for change is essential. Effective engagement lays the foundation for accurate assessment and meaningful intervention, fostering a therapeutic alliance built on trust, empathy, and respect.

Human growth and development provide a framework for understanding how substance use and addiction manifest across the lifespan. CACs and CAPs must recognize the developmental stage of each client, as age, cognitive abilities, emotional maturity, and social context all influence the way addiction is experienced and treated. Adolescents, young adults, and adults each present unique risk factors and treatment considerations. By integrating knowledge of developmental milestones, counselors can tailor interventions, set realistic goals, and anticipate potential challenges in recovery.

The addictive process is complex, involving biological, psychological, and social components. CACs and CAPs explore how substances alter brain chemistry, create dependency, and influence behavior. Understanding the progression of addiction from experimentation to regular use, tolerance, withdrawal, and potential relapse, is essential for effective assessment and treatment planning. Counselors must recognize early warning signs, patterns of use, and co-occurring issues such as mental health disorders or trauma. This knowledge allows for timely interventions and supports long-term recovery strategies.

Client engagement techniques are critical during the initial assessment phase. CACs and CAPs use active listening, motivational interviewing, and open-ended questioning to gather accurate information while maintaining a supportive environment. Establishing clear expectations, explaining confidentiality, and validating client experiences enhance participation and honesty. Engagement strategies also involve

assessing readiness for change, as clients may demonstrate ambivalence or resistance that can impact assessment accuracy and treatment planning.

Assessment instruments provide objective tools to evaluate substance use severity, mental health status, and co-occurring conditions. Common instruments include the ASAM Criteria, DAST, SASSI, AUDIT, and structured intake assessments. CACs use these tools to inform counseling strategies, monitor progress, and identify risk factors. CAPs incorporate clinical judgment to interpret results, integrate findings with medical and psychiatric data, and develop comprehensive treatment plans. Training in these instruments ensures consistent, ethical, and effective assessment practices.

Case study analysis is a practical component of clinical evaluation. By reviewing real or simulated client cases, CACs and CAPs practice integrating client history, assessment findings, and clinical observations into a coherent understanding of client needs. This process allows counselors to identify intervention priorities, develop treatment strategies, and anticipate challenges. Case study discussions foster critical thinking, collaboration, and professional growth, bridging the gap between theory and applied practice.

Documentation of the clinical evaluation is essential for continuity of care, ethical compliance, and professional accountability. CACs record client interactions, assessment results, and initial treatment plans, while CAPs provide more detailed clinical notes and interpretive summaries. Accurate documentation supports supervision, informs ongoing treatment, and ensures adherence to regulatory standards.

ADDICTION COUNSELING

Addiction is not merely a pattern of repeated substance use, it is a complex biopsychosocial condition influenced by biological, psychological, and social factors. Biological contributors include genetics, brain chemistry, and the physiological effects of substances on the nervous system. Psychological factors encompass personality traits, coping strategies, trauma history, and pre-existing mental health disorders. Social influences include family dynamics, peer relationships, cultural expectations, and socioeconomic conditions. Recognizing these interactions enables counselors to approach treatment holistically, rather than addressing symptoms in isolation.

Case Study Example:
A 22-year-old client presents with alcohol use disorder. He has a family history of addiction, experiences anxiety and mild depression, and recently lost his job. A counselor considering biological, psychological, and social factors will identify multiple points of intervention: therapy for mental health, vocational support, and family involvement in treatment planning.

Reflective Exercise:
Write 3–4 sentences reflecting on how each domain, biological, psychological, social—may influence a client you have worked with or studied in a case scenario.

Discussion Prompt:
How does a multidimensional understanding of addiction change the approach to counseling compared to viewing it as only a behavioral problem?

2. Human Growth and Development

Clients enter treatment with unique developmental considerations. Adolescents may face identity formation challenges and peer pressures. Adults often manage complex responsibilities, long-standing behavioral patterns, and relationship stressors. Older

adults may struggle with isolation, chronic illnesses, or cognitive decline. Understanding these stages allows counselors to tailor interventions that are age-appropriate, culturally sensitive, and contextually relevant.

Example:
An adolescent client experimenting with substances might benefit from structured peer-support groups and family counseling, while an adult client may require strategies for workplace stress management alongside therapy.

Reflective Exercise:
Identify the developmental stage of a client and describe how it might influence their motivation, engagement, and treatment goals.

Discussion Prompt:
What are the key differences in engagement strategies between adolescents and adults in addiction counseling?

3. Client Engagement and Therapeutic Alliance

Engagement begins with trust, empathy, and nonjudgmental communication. Techniques such as motivational interviewing, reflective listening, and validation foster participation and readiness for change. Counselors must also recognize transference and countertransference, ensuring personal biases do not interfere with treatment. Effective engagement directly impacts client adherence and long-term recovery outcomes.

Example:
Using motivational interviewing, a counselor helps a resistant client explore ambivalence toward quitting opioids. The counselor listens actively, reflects statements without judgment, and guides the client to articulate personal reasons for change.

Reflective Exercise:
Describe a time when engagement techniques could have altered a client's willingness to participate. Write 3–4 sentences.

Discussion Prompt:
How do transference and countertransference affect client engagement, and what strategies can counselors use to manage them?

4. Assessment Tools and Techniques

Accurate assessment is essential for individualized treatment planning. Tools include:

- PHQ-9: Measures depression severity
- GAD-7: Screens for anxiety
- PTSD Checklist: Identifies trauma-related symptoms
- ACE Questionnaire: Evaluates adverse childhood experiences
- AUDIT & DAST: Screens for alcohol and drug use severity

Tools Explained

1. PHQ-9 (Patient Health Questionnaire-9):
The PHQ-9 is a self-report tool used to assess the severity of depression. It consists of nine items that reflect the core symptoms of depression, including mood, sleep, appetite, and concentration. Scores help clinicians determine the level of depressive symptoms and guide treatment planning, including the need for therapy, monitoring, or referral for psychiatric care.

2. GAD-7 (Generalized Anxiety Disorder-7):
The GAD-7 is a brief screening tool for anxiety symptoms, focusing on worry, nervousness, and tension over the past two weeks. It helps identify clients who may have generalized anxiety disorder and measures severity. This information assists in designing interventions that address both coping strategies and potential comorbid conditions.

3. PTSD Checklist (PCL):
The PTSD Checklist is used to identify trauma-related symptoms and assess their severity. Clients rate symptoms such as intrusive thoughts, avoidance, and hyperarousal. Results support treatment planning for trauma-informed care, including therapy approaches like cognitive processing therapy or EMDR.

4. ACE Questionnaire (Adverse Childhood Experiences):

The ACE Questionnaire evaluates exposure to adverse experiences during childhood, such as abuse, neglect, or household dysfunction. Higher scores indicate greater risk for long-term physical, mental, and behavioral health issues. Understanding ACE scores helps clinicians develop preventative strategies and trauma-informed treatment plans.

5. AUDIT (Alcohol Use Disorders Identification Test):
The AUDIT is a screening tool designed to assess alcohol consumption, drinking behaviors, and related problems. It identifies risky or harmful drinking patterns and can guide decisions regarding intervention, brief counseling, or referral for specialized treatment.

6. DAST (Drug Abuse Screening Test):
The DAST screens for drug use severity and potential substance-related problems, excluding alcohol. It helps determine the level of substance involvement and guides treatment planning, including the intensity of counseling or referral for specialized addiction services.

Counselors combine these tools with clinical interviews, observations, and collateral information to form a comprehensive picture of the client's needs.

Example:
A counselor administering the PHQ-9 to a client who reports frequent sadness and low energy can identify major depressive symptoms and integrate mental health treatment alongside addiction interventions.

Reflective Exercise:
Select one assessment tool and explain how its results would inform treatment planning. Write 3–4 sentences.

Example 1 – PHQ-9:
Using the PHQ-9 to assess depression provides a clear picture of the client's symptom severity. If a client scores in the moderate range, it indicates that they may benefit from both counseling and structured behavioral interventions. This information informs treatment planning by helping prioritize goals, such as improving mood, coping

strategies, and daily functioning. It also guides decisions about monitoring progress or referring to psychiatric care if necessary.

Example 2 – GAD-7:
Administering the GAD-7 allows clinicians to identify the severity of a client's anxiety symptoms. A high score would indicate the need for interventions focused on anxiety management, such as cognitive-behavioral techniques, mindfulness, or stress reduction strategies. This informs treatment planning by highlighting which symptoms require immediate attention and helping establish achievable short-term goals. It also assists in tracking progress over time.

Example 3 – AUDIT:
The AUDIT assesses the level of alcohol use and related risks. If a client's results indicate hazardous drinking, the treatment plan would include brief intervention, education on alcohol effects, and strategies for reducing consumption. This informs treatment planning by tailoring goals to the client's specific drinking patterns and determining the level of support or referral needed for recovery. It also helps prioritize early intervention to prevent progression of alcohol-related problems.

Discussion Prompt:
Why is it important to combine objective assessment tools with clinical judgment in treatment planning?

5. Case Conceptualization and Treatment Planning

Treatment planning involves synthesizing assessment data, developmental considerations, and psychosocial context into actionable strategies. Goals should be SMART: Specific, Measurable, Achievable, Relevant, Time-bound. Interventions must be evidence-based, client-centered, and adaptable to co-occurring disorders. Progress is continuously monitored, with treatment plans updated as needed.

Example:
A client with alcohol use disorder and depression might have a goal to reduce drinking to zero within 90 days, participate in weekly CBT sessions, and attend support groups. The counselor monitors progress weekly and adjusts interventions if setbacks occur.

Reflective Exercise:
Write 3–4 sentences on how you would develop a treatment plan for a client with co-occurring substance use and anxiety disorder.

Discussion Prompt:
Discuss how the SMART framework ensures treatment goals are both practical and measurable.

Example 1 – Smoking Cessation:
Instead of a general goal like "quit smoking," a SMART goal would be: "The client will decrease cigarette use from 15 per day to 5 per day within six weeks, using a nicotine replacement plan and daily tracking." This goal is Specific (reduce cigarette use), Measurable (from 15 to 5 per day), Achievable (gradual reduction), Relevant (supports health improvement), and Time-bound (six weeks). This structure helps the client see progress and stay motivated.

Example 2 – Coping Skills Development:
Rather than saying "improve coping skills," a SMART goal could be: "The client will practice two new stress-reduction techniques, such as deep breathing and journaling, at least five days per week for the next four weeks." The goal is Specific, Measurable (two techniques, five days per week), Achievable, Relevant (supports recovery), and Time-bound (four weeks). Using SMART goals ensures the client has clear, trackable steps to manage triggers effectively.

6. Integration and Ethical Considerations

Ethics are the backbone of counseling. Counselors must uphold confidentiality, professional boundaries, and legal standards while delivering care. Understanding the client's cultural background, past trauma, and potential triggers ensures safe and

effective interventions. Ethical decision-making is woven into every stage of assessment, engagement, and treatment planning.

Example:
Before beginning family therapy, a counselor ensures that each family member consents, understands confidentiality limits, and has a plan for emotional support if sensitive topics arise.

Reflective Exercise:
Identify an ethical dilemma a counselor might face and describe how you would navigate it in 3–4 sentences.

Example 1 – Dual Relationships:
A common ethical dilemma a counselor might face is managing a dual relationship, such as seeing a client outside of therapy in a social setting. I would navigate this by maintaining clear professional boundaries, discussing the situation with the client if necessary, and consulting with a supervisor to ensure ethical standards are upheld. Documenting decisions and actions would also be important for accountability.

Example 2 – Confidentiality vs. Safety:
A counselor may face a dilemma when a client discloses intentions to harm themselves or others. I would prioritize safety by following mandated reporting laws, while explaining confidentiality limits to the client and providing support and resources. Using clinical judgment and supervision helps ensure ethical and legal responsibilities are met.

Discussion Prompt:
How do cultural competence and trauma-informed approaches intersect with ethical counseling practices?

Example 1 – Respecting Cultural Beliefs in Treatment:
Cultural competence ensures counselors understand and respect a client's cultural values, beliefs, and traditions. When combined with trauma-informed care, this approach helps avoid retraumatization and supports ethical practice by honoring client autonomy and identity. For instance, incorporating culturally relevant coping strategies

or rituals into a treatment plan demonstrates both ethical responsibility and sensitivity to trauma.

Example 2 – Communication and Language Considerations:
Using trauma-informed and culturally competent approaches means adapting communication to the client's language and cultural norms. Ethically, counselors are responsible for ensuring clients understand their treatment and can make informed decisions. For example, providing a translator or culturally adapted materials ensures informed consent and supports ethical engagement.

Example 3 – Recognizing Historical Trauma:
Counselors working with clients from marginalized or historically oppressed communities must recognize the impact of intergenerational trauma. Ethically, integrating this understanding into treatment prevents harm and promotes client trust. For example, addressing both substance use and trauma triggers while respecting cultural context aligns with trauma-informed care and ethical standards of beneficence and nonmaleficence.

REFLECTIVE EXERCISE ANSWERS EXAMPLES

1. Write 3–4 sentences reflecting on how each domain, biological, psychological, social may influence a client you have worked with or studied in a case scenario.
Biological factors such as genetics or brain chemistry can influence a client's susceptibility to addiction and response to treatment. Psychological factors, including coping skills, trauma history, and mental health conditions, affect how the client perceives challenges and engages in therapy. Social factors such as family support, peer influence, and socioeconomic conditions can either support or hinder recovery. Understanding all three domains helps counselors develop a holistic treatment plan that addresses the client's unique needs.

2. Identify the developmental stages of a client and describe how it might influence their motivation, engagement, and treatment goals.
A client in adolescence may be motivated by peer acceptance and require structured, interactive sessions, while adults may prioritize practical outcomes that affect their work or family life. Developmental stage shapes cognitive maturity, emotional regulation, and coping strategies, which in turn influence engagement in therapy. Tailoring treatment goals to align with the client's stage ensures interventions are relevant, realistic, and attainable. Recognizing developmental needs improves participation and fosters a supportive therapeutic environment.

3. Describe a time when engagement techniques could have altered a client's willingness to participate. Write 3–4 sentences.
In a case where a client was resistant to group therapy, using motivational interviewing techniques could have encouraged self-reflection on the benefits of participation. Establishing rapport and exploring ambivalence allowed the client to feel heard and understood. Adjusting engagement strategies to the client's readiness to change increased willingness to attend sessions. This demonstrates the importance of individualized approaches in fostering active participation.

4. Select one assessment tool and explain how results would inform treatment planning. Write 3–4 sentences.
Using the PHQ-9 to assess depression provides a measurable score of symptom severity. High scores may indicate the need for concurrent mental health interventions alongside addiction treatment. The results guide counselors in adjusting session focus, selecting appropriate therapy modalities, and monitoring progress over time. Incorporating the tool ensures data-driven decisions that improve treatment outcomes.

5. Write 3–4 sentences on how you would develop a treatment plan for a client with co-occurring substance use and anxiety disorder.
I would begin by conducting a comprehensive assessment to identify the severity and triggers of both the substance use and anxiety symptoms. Goals would be integrated, addressing substance abstinence while implementing anxiety management strategies such as CBT or relaxation techniques. Interventions would be sequenced to support stabilization of symptoms, increase coping skills, and promote engagement in therapy. Progress would be monitored through both objective measures and client self-report to ensure the plan remains effective and client-centered.

6. Identify an ethical dilemma a counselor might face and describe how you would navigate it in 3–4 sentences.
An ethical dilemma may arise when a client discloses intent to harm themselves or others. This needs to following legal and professional guidelines, such as mandatory reporting and implementing safety planning. Maintaining confidentiality while protecting the client and others requires careful communication and documentation. Consulting with supervisors or ethics boards ensures that decisions are both ethical and in the client's best interest.

DISCUSSION PROMPT ANSWERS EXAMPLES

1. How does a multidimensional understanding of addiction change the approach to counseling compared with viewing it only as a behavioral problem?
A multidimensional understanding of addiction considers biological, psychological, and social factors rather than just behavior. Counselors who adopt this approach address not only the substance use but also co-occurring mental health conditions, trauma history, social support, and environmental influences. This allows for integrated treatment plans that target multiple contributing factors, rather than focusing solely on

stopping the behavior. Clients benefit from more holistic care, which can increase engagement and improve outcomes.

2. What are the key differences in engagement strategies between adolescence and adult addiction counseling?
Adolescents often respond better to structured, interactive activities and peer-group interventions, as they are heavily influenced by social factors and identity development. Family involvement and parental guidance are also critical for adolescents. Adults may require flexible scheduling, skill-building for work and family responsibilities, and practical problem-solving strategies. Tailoring engagement strategies to developmental stage ensures that interventions are relevant, respectful, and effective for each client's specific needs.

3. How does countertransference or transference affect client engagement, and what strategies can a counselor use to manage them?
Transference occurs when a client projects feelings onto the counselor, while countertransference occurs when the counselor projects their own feelings onto the client. Both can affect engagement by influencing the therapeutic relationship. Counselors can manage these dynamics through self-awareness, regular supervision, reflective practice, and maintaining professional boundaries. Recognizing these dynamics allows counselors to respond appropriately, maintain trust, and ensure interventions remain client-centered.

4. Why is it important to combine objective assessment tools with clinical judgment in treatment planning?
Objective tools provide measurable data on symptoms, severity, and functioning, which is valuable for tracking progress. However, these tools do not capture the client's full context, such as cultural, social, or trauma-related factors. Combining objective tools with clinical judgment allows counselors to interpret results accurately and develop individualized treatment plans that are both evidence-based and client-specific, improving the likelihood of successful outcomes.

5. Discuss how the SMART framework in insurance treatment goals ensures that goals are both practical and measurable.
The SMART framework (Specific, Measurable, Achievable, Relevant, Time-bound) ensures that treatment goals are clear, realistic, and trackable. Specific goals define exactly what the client will achieve. Measurable goals allow counselors to track progress objectively. Achievable goals are realistic for the client's abilities and resources. Relevant goals align with the client's needs and recovery objectives. Time-bound goals include clear deadlines, which provide motivation and accountability. Using SMART

goals improves treatment planning and helps demonstrate measurable progress for both clients and insurance purposes.

6. How do cultural competence and trauma-informed approaches intersect with ethical counseling practices?
Cultural competence and trauma-informed care require counselors to recognize and respect the client's cultural background, beliefs, and experiences with trauma. Ethical practice involves providing care that is nonjudgmental, safe, and client-centered. By integrating these approaches, counselors ensure interventions are sensitive to the client's context, avoid retraumatization, and maintain professional boundaries. This intersection supports ethical responsibility, promotes client trust, and enhances the effectiveness of treatment outcomes.

Week 1: Clinical Evaluation I (25 hours)
Module Overview

This module introduces trainees to the foundational concepts of addiction, human growth and development, and the clinical evaluation process. Trainees will learn how to identify addiction patterns, engage clients professionally, and begin practicing structured intake assessments.

Learning Objectives

By the end of this module, trainees will be able to:

1. Explain stages of human growth and development relevant to addiction assessment (infancy, childhood, adolescence, adulthood, older adulthood). Include cognitive, emotional, and social development markers.
2. Describe the addictive process, including initiation, escalation, maintenance, and relapse cycles.
3. Identify the signs and symptoms of addiction across various populations (e.g., youth, adults, elderly).
4. Demonstrate client engagement strategies that build rapport and trust during intake.
5. Apply basic intake assessment techniques using structured questions and observations.
6. Document findings accurately using standard intake forms and notes.

Instructor Notes for Teaching

- Human Growth & Development: Provide a visual chart showing age ranges, cognitive/emotional development, and typical risk behaviors. Use examples of how addiction can present at each stage.
- Addictive Process: Explain using a cycle diagram: initiation → experimentation → regular use → abuse → dependence → relapse. Provide real-world examples from clinical experience.
- Client Engagement: Emphasize tone, body language, and open-ended questioning. Discuss cultural and ethical considerations.
- Assessment Instruments: Introduce basic screening tools (e.g., ASAM criteria overview, basic structured interview questions) without going into full scoring.

Topics & Teaching Content

1. Introduction to Addiction

- Definition of substance use disorder and behavioral addictions
- Common signs and symptoms
- Differentiation between casual use and problem use

2. Human Growth & Development Stages

- Infancy and early childhood: attachment, emotional regulation, early risk behaviors
- Adolescence: peer influence, identity formation, experimental substance use
- Adulthood: coping mechanisms, stress factors, co-occurring disorders
- Older adulthood: late-onset addictions, health complications

3. The Addictive Process

- Initiation: first exposure
- Escalation: increasing frequency/intensity
- Maintenance: dependence and routine use
- Relapse: triggers, prevention strategies

4. Client Engagement & Rapport-Building

- Active listening and empathy
- Open-ended questions
- Non-verbal cues and professional demeanor
- Cultural sensitivity

5. Introduction to Intake Assessment

- Structured interviews
- Gathering psychosocial, medical, and substance use history
- Observational skills: appearance, behavior, affect

6. Documentation Basics

- Writing clear intake notes
- Recording objective vs. subjective information
- Maintaining confidentiality

1. Introduction to Addiction

This section provides learners with foundational knowledge about substance use disorders (SUDs) and behavioral addictions. Begin by defining SUDs as chronic, relapsing conditions characterized by compulsive use despite negative consequences, and behavioral addictions as compulsive engagement in non-substance behaviors (e.g., gambling, gaming). Highlight common signs and symptoms, such as impaired control, cravings, withdrawal, tolerance, and functional impairment. Emphasize the difference between casual use and problem use using practical examples—e.g., occasional social drinking versus daily drinking leading to health, legal, or social problems. Discussing this helps learners identify at-risk clients early and understand the spectrum of addictive behaviors.

Example: A college student drinks at parties weekly (casual use), versus another student drinking daily and missing classes (problem use).